

TSSA Return to Play Form (Concussion)

This form is adapted from the Acute Concussion Evaluation Plan on the Centers for Disease and Control and Prevention (CDC) website. All medical providers are encouraged to review this site if they have questions regarding the latest information on the evaluation and care of the youth athlete following a concussion injury (www.cdc.gov/injury). *Please initial any recommendations selected*.

Print Athletes Name:		Date of Injury:	Date of Birth:
This return to play plan is based on today's evaluation. Care plane completed by:		Date of Evaluation: Return to this Office (Date): Return to School on (Date):	
Return to Sports: 1. Athletes should not return to practice or competition the same day their head injury occurred. 2. Athletes should never return to play or competition if they still have ANY symptoms. 3. Athletes, be sure your coach or team manager is aware of your injury, symptoms, and has the contact information for the treating health care provider.			
The following are return to sports recommendations at the present time:			
PHYSICAL EDUCATION:	Do NOT return to PE clas	s at this timeMay r	eturn to PE class
SPORTS (SOCCER):	health care professional (May be advanced back to care professional	soccer practice or competition	ation with the treating health
-OR-		on in all activities without res	striction
Treating Health Care Provider Information (Please Print/Stamp)			
Please check: Medical Doctor (M.D.) Provider's Name: Provider's Signature:	Osteopathic Physician (D.O.) Provid Office	Clinical Neuropsycholer's Office Phone: Address:	

Gradual Return to Play of Concussed Athlete

Baseline: Athletes should not have any concussion symptoms. Athletes should only progress to the next step if they do not have any symptoms at the current step.

- **Step1:** Begin with light aerobic exercise only to increase an athlete's heart rate. This means about 10 minutes on an exercise bike, walking, or light jogging. No weightlifting at this point.
- Step 2: Continue with activities to increase an athlete's heart rate with body or head movement. This includes moderate jogging, brief running, moderate-intensity stationary biking, moderate-intensity weightlifting (reduced time and/or reduced weight from their typical routine).
- *Step 3:* Add heavy non-contact physical activity such as sprinting/running, high-intensity stationary biking, and regular weightlifting routine and/or non-contact sport-specific drills (in three planes of movement).
- Step 4: Athlete may return to practice and full contact (if appropriate for the sport) in controlled practice.
- Step 5: Athlete may return to competition.

If an athlete's symptoms come back or they get new symptoms when becoming more active at any step, this is a sign that the athlete is pushing too hard. The athlete should stop these activities and the athlete's health care provider should be contacted. After more rest and no concussion symptoms, the athlete should begin at the previous step.